Author Index

Abramovich-Bar, S., see Woolfson-Bartfeld, D. 305
Abu-Lafi, S., see Levin, S. 285
Aguilar, M. I., see Hodder, A. N. 317
Amari, J. V.
——, Brown, P. R., Grill, C. M. and Turcotte, J. G.
Isolation and purification of lecithin by preparative high-performance liquid chromatography 219
Bamberger, Y., see Woolfson-Bartfeld, D. 305
Bartle, K. D.
——, Clifford, A. A., Jafar, S. A., Kinthinji, J. P. and Shilstone, G. F.
Use of chromatographic retention measurements to obtain solubilities in a liquid or supercritical fluid mobile phase 459
Batten, C. F., see Wentworth, W. E. 87
Bayer, E., see Ilg, M. 263
Beckett, R.
—, Hotchin, D. M. and Hart, B. T.
Use of field-flow fractionation to study pollutant–colloid interactions 435
Bertsch, W., see Brill, J. H. 95
Boone, L. S., see Kirkland, J. J. 377
Brill, J. J.
—— and Bertsch, W.
Comparison of cuticular hydrocarbon profiles of fire ants *Solenopsis richteri* from the same colony, using capillary column gas chromatography with pattern recognition 95
Brown, P. R., see Amari, J. V. 219
Bruin, G. J. M.
——, Tock, P. P. H., Kraak, J. C. and Poppe, H.
Electrically driven open-tubular liquid chromatography 557
Bruner, F.
——, Crescentini, G., Mangani, F. and Lattanzi, L.
Capillary gas chromatography with graphitized carbon black 123
Caldwell, C. D., see Li, J. 361
Carr, P. W., see Li, J. 103
Chang, H.-C. K.
—— and Taylor, L. T.
Use of sulfur chemiluminescence detection after supercritical fluid chromatography 491
Chen, E. C. M., see Wentworth, W. E. 87
Clifford, A. A., see Bartle, K. D. 459
Cramers, C. A., see Van Es, A. J. 143
Crescentini, G., see Bruner, F. 123
Dallas, A. J., see Li, J. 103
Davis, J. M.
Influence of thermal variation of diffusion coefficient on non-equilibrium plate height in capillary zone electrophoresis 521
D'sa, E. D., see Wentworth, W. E. 87
Engewald, W., see Maurer, T. 77
Es, A. J. van, see Van Es, A. J. 143
Friedrich, J. P., see King, J. W. 449
Giddings, J. C., see Moon, M. H. 423
Goebel, L. K., see Rasmussen, H. T. 549
Golay, M. J. E., see Van Es, A. J. 143
Golovnya, R. V.
—— and Polanuer, B. M.
Comparison of methods for the determination of the polarity and selectivity of stationary phases in gas chromatography from a thermodynamic point of view 51
Golshan-Shirazi, S.
——, and Guiochon, G.
Optimization of the experimental conditions in preparative liquid chromatography with touching bands 229
Grill, C. M., see Amari, J. V. 219
Grushka, E., see Woolfson-Bartfeld, D. 305
Guiochon, G., see Golshan-Shirazi, S. 229
Hansen, M. E.
—— and Short, D. C.
Optimization study of octane-in-water emulsions by sedimentation field-flow fractionation 333
Hanson, M.
——, Unger, K. K. and Schomburg, G.
Non-porous polybutadiene-coated silicas as stationary phases in reversed-phase chromatography 269
Haraguchi, H., see Takeuchi, T. 257
Hart, B. T., see Beckett, R. 435
Hearn, M. T. W., see Hodder, A. N. 317
Hevesi, T.
——, Krupčík, J. and Sandra, P.
Use of chromatographic models for computerized optimization of coupling-point pressure in dual-column gas chromatography 161
Hodder, A. N.
——, Machin, K. J., Aguilar, M. I. and Hearn, M. T. W.
High-performance liquid chromatography of amino acids, peptides and proteins. C. Characterisation of coulombic interactive regions on hen lysozyme by high-performance liquid anion-exchange chromatography and computer graphic analysis 317
Hotchin, D. M., see Beckett, R. 435

Housaki, T., see Nishikida, K. 209
Hu, W., see Takeuchi, T. 257
Huang, E. C., see Sumpter, S. R. 503
Ilg, M.
——-, Maier-Rosenkranz, J., Müller, W. and Bayer, E.
Magnetic resonance imaging in reversed-phase liquid chromatography 263
Ishii, D., see Takeuchi, T. 257
Itoh, K., see Jinno, K. 193
Jafar, S. A., see Bartle, K. D. 459
Jinno, K.
——, Yamamoto, K., Nagashima, H., Ueda, T. and Itoh, K.
Silicas chemically bonded with multidentate phenyl groups as stationary phases in reversed-phase liquid chromatography used for non-planarity recognition of polycyclic aromatic hydrocarbons 193
Kaljurand, M., see Küllik, E. 175
Karaiskakis, G., see Koliadima, A. 345
King, J. W.
—— and Friedrich, J. P.
Quantitative correlations between solute molecular structure and solubility in supercritical fluids 449
Kinoshita, T., see Nishikida, K. 209
Kirkland, J. J.
——, Boone, L. S. and Yau, W. W.
Retention effects in thermal field-flow fractionation 377
Kithinji, J. P., see Bartle, K. D. 459
Koch, L.
——, Koch, T. and Widmer, H. M.
Sedimentation field-flow fractionation for pigment quality assessment 395
Koch, T., see Koch, L. 395
Koliadima, A.
—— and Karaiskakis, G.
Potential-barrier field-flow fractionation, a versatile new separation method 345
Kraak, J. C., see Bruin, G. J. M. 557
Krupčík, J., see Hevesi, T. 161
Küllik, E.
—— and Kaljurand, M.
Study of rapid reaction kinetics by computerized gas chromatography with stroboscopic sampling 175
Lattanzi, L., see Bruner, F. 123
Leem, M. L., see Sumpter, S. R. 503
Levin, S.
—— and Abu-Lafi, S.
Competition between penylalanine and acetic acid in a chromatographic column as indicated by their adsorption isotherms 285
Levy, S., see Woolfson-Bartfeld, D. 305
Li, J.
——, Caldwell, K. D. and Mächtle, W.
Particle characterization in centrifugal fields. Comparison between ultracentrifugation and sedimentation field-flow fractionation 361
——, Dallas, A. J. and Carr, P. W.
Empirical scheme for the classification of gas chromatographic stationary phases based on solvatochromic linear solvation energy relationships 103Luffer, D. R.
Luffer, D. R.
—— and Novotny, M.
Element-selective detection after supercritical fluid chromatography by means of a surfatron plasma in the near-infrared spectral region 477
Machin, K. J., see Hodder, A. N. 317
McNair, H. M., see Rasmussen, H. T. 549
Mächtle, W., see Li, J. 361
Magnico, P.
—— and Martin, M.
Dispersion in the interstitial space of packed columns 31
Maier-Rosenkranz, J., see Ilg, M. 263
Mangani, F., see Bruner, F. 123
Markides, K. E., see Sumpter, S. R. 503
Martin, M., see Magnico, P. 31
Martire, D. E., see Poe, D. P. 3
Maurer, T.
——, Engewald, W. and Steinborn, A.
Enhanced possibilities for identification using series-coupled capillary gas chromatographic columns. II. Retention indices as an identification tool in selectivity tuning 77
Moon, M. H.
——, Myers, M. N. and Giddings, J. C.
Evaluation of pinched inlet channel for stopless flow injection in steric field-flow fractionation 423
Morimoto, M., see Nishikida, K. 209
Müller, W., see Ilg, M. 263
Myers, M. N., see Moon, M. H. 423
Nagashima, H., see Jinno, K. 193
Nishikida, K.
——, Housaki, T., Morimoto, M. and Kinoshita, T.
Gel permeation chromatography–Fourier transform infrared study of some synthetic polymers. II. Instrumentation for the characterization of polyethylene 209
Novotny, M., see Luffer, D. R. 477
Oguchi, R., see Wylie, P. L. 131
Poe, D. P.
—— and Martire, D. E.
Plate height theory for compressible mobile phase fluids and its application to gas, liquid and supercritical fluid chromatography 3

Polanuer, B. M., see Golovnya, R. V. 51
Poppe, H., see Bruin, G. J.M. 557
Rasmussen, H. T.
——, Goebel, L. K. and McNair, H. M.
Micellar electrokinetic chromatography employing sodium alkyl sulfates and Brij 35® 549
Rijks, J. A., see Van Es, A. J. 143
Sandra, P., see Hevesi, T. 161
Schimpf, M. E.
Characterization of polymers by thermal field-flow fractionation 405
Schomburg, G., see Hanson, M. 269
Scott, R. P. W.
Design of liquid chromatography capillary columns 297
Shilstone, G. F., see Bartle, K. D. 459
Short, D. C., see Hansen, M. E. 333
Smith, R. D., see Yonker, C. R. 573
Springston, S. R.
Cryogenic-focusing, ohmically heated on-column trap for capillary gas chromatography 67
Steinborn, A., see Maurer, T. 77
Sumpter, S. R.
——, Woolley, C. L., Huang, E. C., Markides, K. E. and Lee, M. L.
Static coating of 5 to 50 μm I.D. capillary columns for open tubular column chromatography 503
Takeuchi, T.
——, Hu, W., Haraguchi, H. and Ishii, D.
Evaluation of the stability of polymer-coated silica-based packing materials for high-performance liquid chromatography 257
Taylor, L. T., see Chang, H.-C. K. 491
Tock, P. P. H., see Bruin, G. J. M. 557
Turcotte, J. G., see Amari, J. V. 219
Uden, P. C., see Wang, T. 185
Ueda, T., see Jinno, K. 193
Unger, K. K., see Hanson, M. 269
Van Es, A. J.
——, Rijks, J. A., Cramers, C. A. and Golay, M. J. E.
Turbulent flow in capillary gas chromatography—evaluation of a theoretical concept by Golay 143
Wang, T.
—— and Uden, P. C.
High-performance liquid chromatographic study of ligand-exchange reactions of fluorinated metal β-diketone chelates 185
Wentworth, W. E.
——, D'sa, E. D., Batten, C. F. and Chen, E. C. M.
Oxygen enhancement of thermal electron capture in a non-radioactive discharge source of a quadrupole mass spectrometer 87
Widmer, H. M., see Koch, L. 395
Woolfson-Bartfeld, D.
——, Grushka, E., Abramovich-Bar, S., Levy, S. and Bamberger, Y.
Reversed-phase ion-pair chromatography with indirect photometric detection of inorganic anions from residues of low explosives 305
Woolley, C. L., see Sumpter, S. R. 503
Wylie, P. L.
—— and Oguchi, R.
Pesticide analysis by gas chromatography with a novel atomic emission detector 131
Yamamoto, K., see Jinno, K. 193
Yau, W. W., see Kirkland, J. J. 377
Yonker, C. R.
—— and Smit, R. D.
High-pressure and supercritical capillary electrophoresis 573